



Luciano Casimiro

MAY 4, 2010 - SEP 7, 2010



Scan to Visit

STORYLAND
FUNERAL HOME

Table of Contents

Obituary	Page 3
Events	Page 5



Luciano Casimiro

MAY 4, 2010 - SEP 7, 2010

Flowering plants usually face selective pressure to optimise the transfer of their pollen, and this is typically reflected in the morphology of the flowers and the behaviour of the plants. Pollen may be transferred between plants via a number of 'vectors'. Some plants make use of abiotic vectors — namely wind (anemophily) or, much less commonly, water (hydrophily). Others use biotic vectors including insects (entomophily), birds (ornithophily), bats (chiropterophily) or other animals. Some plants make use of multiple vectors, but many are highly specialised. Cleistogamous flowers are self pollinated, after which they may or may not open. Many *Viola* and some *Salvia* species are known to have these types of flowers. The flowers of plants that make use of biotic pollen vectors commonly have glands called nectaries that act as an incentive for animals to visit the flower. Some flowers have patterns, called nectar guides, that show pollinators where to look for nectar. Flowers also attract pollinators by scent and color. Still other flowers use mimicry to attract pollinators. Some species of orchids, for example, produce flowers resembling female bees in color, shape, and scent. Flowers are also specialized in shape and have an arrangement of the stamens that ensures that pollen grains are transferred to the bodies of the pollinator when it lands in search of its attractant (such as nectar, pollen, or a mate). In pursuing this attractant from many flowers of the same species, the pollinator transfers pollen to the stigmas—arranged with equally pointed precision—of all of the flowers it visits. Anemophilous flowers use the wind to move pollen from one flower to the next. Examples include grasses, birch trees, ragweed and maples. They have no need to attract pollinators and therefore tend not to be "showy" flowers. Male and female reproductive organs are generally found in separate flowers, the male flowers having a number of long filaments terminating in exposed stamens, and the female flowers having long, feather-like stigmas. Whereas the pollen of animal-pollinated flowers



Obituary

Luciano Casimiro

MAY 4, 2010 - SEP 7, 2010

tends to be large-grained, sticky, and rich in protein (another "reward" for pollinators), anemophilous flower pollen is usually small-grained, very light, and of little nutritional value to animals.



Events


Luciano Casimiro


MAY 4, 2010 - SEP 7, 2010

BBQ

 **Friday**, September 3, 2010

 8:00 AM - 10:00 AM ET

 **Funeral Las Vegas**
1593 N River Rd, St Clair MI 48079





Memories only last if you share them

Join us in honoring Luciano by contributing to a collection of shared memories.



Scan to Visit